

## Tilburg University

### Association schemes

Haemers, W.H.; Brouwer, A.E.

*Publication date:*  
1987

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

Haemers, W. H., & Brouwer, A. E. (1987). *Association schemes*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 258). Unknown Publisher.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

7626  
387  
CBM  
R  
7626  
1987  
258

UNIVERSITY  
LIJKE  
UNIVERSITEIT  
BRABANT

POSTBOX 90153  
5000 LE TILBURG  
THE NETHERLANDS



DEPARTMENT OF ECONOMICS  
RESEARCH MEMORANDUM



## ASSOCIATION SCHEMES

W.H. Haemers

A.E. Brouwer

FEW 258



ASSOCIATION SCHEMES

A.E. Brouwer,  
Technological University Eindhoven

W.H. Haemers,  
Tilburg University

This paper gives a brief survey of the theory of Association Schemes. It is destined to become a chapter of the Handbook of Combinatorics.

Table of contents

1. Introduction	1
2. Strongly Regular Graphs	2
3. Association Schemes	4
4. Applications	8

## Association schemes.

*A.E. Brouwer & W.H. Haemers*

### 1. Introduction.

Association schemes are by far the most important unifying concept in algebraic combinatorics. They provide a common view point for the treatment of problems in several fields, such as coding theory, design theory, algebraic graph theory and finite group theory.

Roughly, an association scheme is a very regular partition of the edge-set of a complete graph. Each of the partition classes defines a graph, and the adjacency matrices of these graphs (together with the identity matrix) form the basis of an algebra (known as the Bose-Mesner algebra of the scheme). But since this basis consists of 0-1 matrices, we see that the algebra is not only closed under matrix multiplication, but also under componentwise (Hadamard, Schur) multiplication. The interplay between these two algebra structures on the Bose-Mesner algebra yields strong information on the structure and parameters of an association scheme.

The relation of the theory of association schemes to the various fields varies. Finite group theory mainly serves as a source of examples - any generously transitive permutation representation of a group yields an association scheme -, but also as a source of inspiration - many group-theoretic concepts and results can be mimicked in the (more general) setting of association schemes. On the other hand, in combinatorics association schemes form a tool. Using inequalities for the parameters of an association scheme, one finds upper bounds for the size of error-correcting codes, and lower bounds for the number of blocks of a  $t$ -design. Many uniqueness proofs for combinatorial structures depend crucially on the additional information one gets in case such inequalities hold with equality.

The graphs of the partition classes of an association scheme are very special. For instance distance-regular graphs and Kneser graphs are of this type. For these graphs the spectrum of the adjacency matrix can be computed from the parameters of the association scheme. Thus results about the eigenvalues of graphs (cf. Godsil's chapter) can be applied.

An association scheme with just two classes is the same as a pair of complementary strongly regular graphs. Because strongly regular graphs are themselves important combinatorial objects and because their treatment forms a good introduction to the more general theory, we shall start with a section on strongly regular graphs. In the subsequent section we treat the elements of the general theory of association schemes: the adjacency matrices, the Bose-Mesner algebra, the eigenvalues, the Krein parameters, the absolute bound and Delsarte's inequality on subsets of an association scheme. The last section is devoted to some special association schemes and the significance to other fields of combinatorics, as indicated above.

Association schemes were introduced by BOSE & SHIMAMOTO [10] and BOSE & MESNER [9] defined the algebra that bears their name. DELSARTE [17] found the linear programming bound, studied  $P$ - and  $Q$ -polynomial schemes, and applied the resulting theory to codes and designs. HIGMAN [29] introduced the more general concept of coherent configuration to study permutation representations of finite groups. There is much literature on association schemes and related topics. BANNAI & ITO [3] is the first text book on association schemes. On the more special topic of distance-regular graphs a lot of material can be found in BIGGS [7]; a monograph on this topic by Brouwer, Cohen & Neumaier is under preparation. Some introductory papers on association schemes are DELSARTE [18], GOETHALS [23], chapter 21 of MACWILLIAMS & SLOANE [43], HAEMERS [25], chapter 17 of CAMERON & VAN LINT [15] and SEIDEL [50].



## 2. Strongly regular graphs.

A simple graph of order  $v$  is *strongly regular* with parameters  $v, k, \lambda, \mu$  whenever it is not complete or empty and

- (i) each vertex is adjacent to  $k$  vertices,
- (ii) for each pair of adjacent vertices there are  $\lambda$  vertices adjacent to both,
- (iii) for each pair of non-adjacent vertices there are  $\mu$  vertices adjacent to both.

For example, the pentagon is strongly regular with parameters  $(v, k, \lambda, \mu) = (5, 2, 0, 1)$ . One easily verifies that a graph  $G$  is strongly regular with parameters  $(v, k, \lambda, \mu)$  if and only if its complement  $\bar{G}$  is strongly regular with parameters  $(v, v-k-1, v-2k+\mu-2, v-2k+\lambda)$ . The line graph of the complete graph of order  $m$ , known as the *triangular graph*  $T(m)$ , is strongly regular with parameters  $(\frac{1}{2}m(m-1), 2(m-2), m-2, 4)$ . The complement of  $T(5)$  has parameters  $(10, 3, 0, 1)$ . This is the Petersen graph.

A graph  $G$  satisfying condition (i) is called  $k$ -regular. It is well-known and easily seen that the adjacency matrix of a  $k$ -regular graph has an eigenvalue  $k$  with eigenvector  $j$  (the all-one vector), and that every other eigenvalue  $\rho$  satisfies  $|\rho| \leq k$  (see Godsil's chapter or BIGGS [7]). For convenience we call an eigenvalue *restricted* if it has an eigenvector perpendicular to  $j$ . We let  $I$  and  $J$  denote the identity and all-one matrices, respectively.

**2.1. Theorem.** *For a simple graph  $G$  of order  $v$ , not complete or empty, with adjacency matrix  $A$ , the following are equivalent:*

- (i)  $G$  is strongly regular with parameters  $(v, k, \lambda, \mu)$  for certain integers  $k, \lambda, \mu$ ,
- (ii)  $A^2 = (\lambda - \mu)A + (k - \mu)I + \mu J$  for certain reals  $k, \lambda, \mu$ ,
- (iii)  $A$  has precisely two distinct restricted eigenvalues.

**Proof.** The equation in (ii) can be rewritten as

$$A^2 = kI + \lambda A + \mu(J - I - A).$$

Now (i)  $\Leftrightarrow$  (ii) is obvious. (ii)  $\Rightarrow$  (iii): Let  $\rho$  be a restricted eigenvalue, and  $u$  a corresponding eigenvector perpendicular to  $j$ . Then  $Ju = 0$ . Multiplying the equation in (ii) on the right by  $u$  yields  $\rho^2 = (\lambda - \mu)\rho + (k - \mu)$ . This quadratic equation in  $\rho$  has two distinct solutions. (Indeed,  $(\lambda - \mu)^2 = 4(\mu - k)$  is impossible since  $\mu \leq k$  and  $\lambda \leq k - 1$ .)

(iii)  $\Rightarrow$  (ii): Let  $r$  and  $s$  be the restricted eigenvalues. Then  $(A - rI)(A - sI) = \alpha J$  for some real number  $\alpha$ . So  $A^2$  is a linear combination of  $A$ ,  $I$  and  $J$ . ■

As an application, we show that quasisymmetric block designs give rise to strongly regular graphs. A quasisymmetric design is a  $2-(v, k, \lambda)$  design such that any two blocks meet in either  $x$  or  $y$  points, for certain fixed  $x, y$ . (Cf. the chapter on block designs.) Given this situation, we may define a graph  $G$  on the set of blocks, and call two blocks adjacent when they meet in  $x$  points. Then there exist coefficients  $\alpha_1, \dots, \alpha_7$  such that  $NN^T = \alpha_1 I + \alpha_2 J$ ,  $NJ = \alpha_3 J$ ,  $JN = \alpha_4 J$ ,  $A = \alpha_5 N^T N + \alpha_6 I + \alpha_7 J$ , where  $A$  is the adjacency matrix of the graph  $G$ . (The  $\alpha_i$  can be readily expressed in terms of  $v, k, \lambda, x, y$ .) Then  $G$  is strongly regular by (ii) of the previous theorem. (Indeed, from the equations just given it follows straightforwardly that  $A^2$  can be expressed as a linear combination of  $A$ ,  $I$  and  $J$ .) A large class of quasisymmetric block designs is provided by the  $2-(v, k, \lambda)$  designs with  $\lambda = 1$  (also known as Steiner systems  $S(2, k, v)$ ) - such designs have only two intersection numbers since no two blocks can meet in more than one point. This leads to a substantial family of strongly regular graphs, including the triangular graphs  $T(m)$  (derived from the trivial design consisting of all pairs out of an  $m$ -set).

Another connection between strongly regular graphs and designs is found as follows: Let  $A$  be the adjacency matrix of a strongly regular graph with parameters  $(v, k, \lambda, \mu)$  (i.e., with  $\lambda = \mu$ ; such a graph is sometimes called a  $(v, k, \lambda)$  graph). Then, by (2.1.ii)

$$AA^T = A^2 = (k - \lambda)I + \lambda J,$$

which reflects that  $A$  is the incidence matrix of a square ('symmetric')  $2-(v, k, \lambda)$  design. (And in this way one obtains precisely all square 2-designs possessing a polarity without absolute points.) For instance, the triangular graph  $T(6)$  provides a square  $2-(15, 8, 4)$  design, the complementary design of the design of



points and planes in the projective space  $PG(3,2)$ . Similarly, if  $A$  is the adjacency matrix of a strongly regular graph with parameters  $(v, k, \lambda, \lambda+2)$ , then  $A+I$  is the incidence matrix of a square 2- $(v, k, \lambda)$  design (and in this way one obtains precisely all square 2-designs possessing a polarity with all points absolute).

**2.2. Theorem.** *Let  $G$  be a strongly regular graph with adjacency matrix  $A$  and parameters  $(v, k, \lambda, \mu)$ . Let  $r$  and  $s$  ( $r > s$ ) be the restricted eigenvalues of  $A$  and let  $f, g$  be their respective multiplicities. Then*

- (i)  $k(k-1-\lambda) = \mu(v-k-1)$ ,
- (ii)  $rs = \mu - k, \quad r+s = \lambda - \mu$ ,
- (iii)  $f, g = \frac{1}{2}(v-1) \mp \frac{(r+s)(v-1)+2k}{r-s}$ .

**Proof.** (i) Fix a vertex  $x$  of  $G$ . Let  $\Gamma(x)$  and  $\Delta(x)$  be the sets of vertices adjacent and non-adjacent to  $x$ , respectively. Counting in two ways the number of edges between  $\Gamma(x)$  and  $\Delta(x)$  yields (i). The equations (ii) are direct consequences of (2.1.ii), as we saw in the proof. Formula (iii) follows from  $f+g = v-1$  and  $0 = \text{tr } A = k + fr + gs = k + \frac{1}{2}(r+s)(f+g) + \frac{1}{2}(r-s)(f-g)$ . ■

These relations imply restrictions for the possible values of the parameters. Clearly, the right hand sides of (iii) must be positive integers. These are the so-called *rationality conditions*. They imply that  $r$  and  $s$  must be integers if  $f \neq g$ . As an example of the application of the rationality conditions we can derive the following result due to HOFFMAN & SINGLETON [32].

**2.3. Theorem.** *Suppose  $(v, k, 0, 1)$  is the parameter set of a strongly regular graph. Then  $(v, k) = (5, 2), (10, 3), (50, 7)$  or  $(3250, 57)$ .*

**Proof.** The rationality conditions imply that either  $f = g$ , which leads to  $(v, k) = (5, 2)$ , or  $r-s$  is an integer dividing  $(r+s)(v-1)+2k$ . By use of (2.2.i-ii) we have

$$s = -r-1, \quad k = r^2+r+1, \quad v = r^4+2r^3+3r^2+2r+2,$$

and thus we obtain  $r = 1, 2$  or  $7$ . ■

The first three possibilities are uniquely realized by the pentagon, the Petersen graph and the Hoffman-Singleton graph. For the last case existence is unknown (but see ASCHBACHER [1]).

Except for the rationality conditions, other restrictions on the parameters are known. We mention three of them.

The Krein conditions, due to SCOTT [48], can be stated as follows:

$$\begin{aligned} (r+1)(k+r+2rs) &\leq (k+r)(s+1)^2, \\ (s+1)(k+s+2rs) &\leq (k+s)(r+1)^2. \end{aligned}$$

Seidel's absolute bound (see DELSARTE, GOETHALS & SEIDEL [20], KOORNWINDER [37], SEIDEL [49]) reads

$$\begin{aligned} v &\leq f(f+3), \\ v &\leq g(g+3). \end{aligned}$$

The conference matrix condition, due to BELEVITCH [4] (see also VAN LINT & SEIDEL [41]), states that if  $f = g$ , then  $v$  must be the sum of two squares (such a graph is related to a conference matrix of order  $v+1$ ).

The Krein conditions and the absolute bound are special cases of general inequalities for association schemes - we'll meet them again in the next section; the conference matrix condition is the analogue of the Bruck-Chowla-Ryser theorem for square 2-designs. In BROUWER & VAN LINT [11] one may find a list of all known restrictions; this paper gives a survey of the recent results on strongly regular graphs. It is a sequel to HUBAUT [33]'s earlier survey of constructions. SEIDEL [49] gives a good treatment of the theory. Some other references are BOSE [8], CAMERON [14] and CAMERON & VAN LINT [15].

### 3. Association schemes.

An *association scheme with  $d$  classes* is a finite set  $X$  together with  $d+1$  relations  $R_i$  on  $X$  such that

- (i)  $\{R_0, R_1, \dots, R_d\}$  is a partition of  $X \times X$ ;
- (ii)  $R_0 = \{(x, x) | x \in X\}$ ;
- (iii) if  $(x, y) \in R_i$ , then also  $(y, x) \in R_i$ , for all  $x, y \in X$  and  $i \in \{0, \dots, d\}$ ;
- (iv) for any  $(x, y) \in R_k$  the number  $p_{ij}^k$  of  $z \in X$  with  $(x, z) \in R_i$  and  $(z, y) \in R_j$  depends only on  $i, j$  and  $k$ .

The numbers  $p_{ij}^k$  are called the *intersection numbers* of the association scheme. The above definition is the original definition of BOSE & SHIMAMOTO [10]; it is what DELSARTE [17] calls a symmetric association scheme. In Delsarte's more general definition, (iii) is replaced by:

- (iii') for each  $i \in \{0, \dots, d\}$  there exists a  $j \in \{0, \dots, d\}$  such that  $(x, y) \in R_i$  implies  $(y, x) \in R_j$ ,
- (iii'')  $p_{ij}^k = p_{ji}^k$ , for all  $i, j, k \in \{0, \dots, d\}$ .

D.G. HIGMAN [29, 30, 31] studied the even more general concept of a *coherent configuration*. He requires (i), (iii'), (iv) and

- (ii')  $\{(x, x) | x \in X\}$  is a union of some  $R_j$ .

If (ii) holds, then the coherent configuration is called *homogeneous*. We shall not treat coherent configurations here, but content ourselves with the remark that a coherent configuration with at most 5 classes must be an association scheme (in the sense of Delsarte), see HIGMAN [30].

Define  $n := |X|$ , and  $n_i := p_{ii}^0$ . Clearly, for each  $i \in \{1, \dots, d\}$ ,  $(X, R_i)$  is a simple graph which is regular of degree  $n_i$ .

**3.1. Theorem.** *The intersection numbers of an association scheme satisfy*

- (i)  $p_{0j}^k = \delta_{jk}$ ,  $p_{ij}^0 = \delta_{ij} n_j$ ,  $p_{ij}^k = p_{ji}^k$ ,
- (ii)  $\sum_i p_{ij}^k = n_j$ ,  $\sum_j n_j = n$ ,
- (iii)  $p_{ij}^k n_k = p_{ik}^j n_j$ ,
- (iv)  $\sum_i p_{ij}^l p_{kl}^m = \sum_i p_{ki}^l p_{il}^m$ .

**Proof.** (i)-(iii) are straightforward. The expressions at both sides of (iv) count quadruples  $(w, x, y, z)$  with  $(w, x) \in R_i$ ,  $(x, y) \in R_j$ ,  $(y, z) \in R_k$ , for a fixed pair  $(w, z) \in R_m$ . ■

It is convenient to write the intersection numbers as entries of the so-called *intersection matrices*  $L_0, \dots, L_d$ :

$$(L_i)_{kj} = p_{ij}^k.$$

Note that  $L_0 = I$ . From the definition it is clear that an association scheme with two classes is the same as a pair of complementary strongly regular graphs. If  $(X, R_1)$  is strongly regular with parameters  $(v, k, \lambda, \mu)$ , then the intersection matrices of the scheme are

$$L_1 = \begin{bmatrix} 0 & k & 0 \\ 1 & \lambda & k-\lambda-1 \\ 0 & \mu & k-\mu \end{bmatrix} \quad L_2 = \begin{bmatrix} 0 & 0 & v-k-1 \\ 0 & k-\lambda-1 & v-2k+\lambda \\ 1 & k-\mu & v-2k+\mu-2 \end{bmatrix}.$$

We see that (iii) generalises (2.2.i).

#### The Bose-Mesner algebra.

The relations  $R_i$  of an association scheme are described by their adjacency matrices  $A_i$  of order  $n$  defined by

$$(A_i)_{xy} = \begin{cases} 1 & \text{whenever } (x, y) \in R_i, \\ 0 & \text{otherwise.} \end{cases}$$

In other words,  $A_i$  is the adjacency matrix of the graph  $(X, R_i)$ . In terms of the adjacency matrices, the

axioms (i)-(iv) become

- (i)  $\sum_{i=0}^d A_i = J$ ,
- (ii)  $A_0 = I$ ,
- (iii)  $A_i = A_i^T$ , for all  $i \in \{0, \dots, d\}$ ,
- (iv)  $A_i A_j = \sum_k p_{ij}^k A_k$ , for all  $i, j, k \in \{0, \dots, d\}$ .

From (i) we see that the  $A_i$  are linearly independent, and by use of (ii)-(iv) we see that they generate a commutative  $(d+1)$ -dimensional algebra  $\mathbf{A}$  of symmetric matrices with constant diagonal. This algebra was first studied by BOSE & MESNER [9] and is called the Bose-Mesner algebra of the association scheme.

Since the matrices  $A_i$  commute, they can be diagonalized simultaneously (see MARCUS & MINC [44]), that is, there exist a matrix  $S$  such that for each  $A \in \mathbf{A}$ ,  $S^{-1}AS$  is a diagonal matrix. Therefore  $\mathbf{A}$  is semisimple and has a unique basis of minimal idempotents  $E_0, \dots, E_n$  (see BURROW [12]). These are matrices satisfying

$$\begin{aligned} E_i E_j &= \delta_{ij} E_i, \\ \sum_{i=0}^d E_i &= I. \end{aligned}$$

The matrix  $\frac{1}{n}J$  is a minimal idempotent (idempotent is clear, and minimal follows since  $\text{rk } J = 1$ ). We shall take  $E_0 = \frac{1}{n}J$ . Let  $P$  and  $\frac{1}{n}Q$  be the matrices relating our two bases for  $\mathbf{A}$ :

$$\begin{aligned} A_j &= \sum_{i=0}^d P_{ij} E_i, \\ E_j &= \frac{1}{n} \sum_{i=0}^d Q_{ij} A_i. \end{aligned}$$

Then clearly

$$PQ = QP = nI.$$

It also follows that

$$A_j E_i = P_{ij} E_i,$$

which shows that the  $P_{ij}$  are the eigenvalues of  $A_j$  and that the columns of  $E_i$  are the corresponding eigenvectors. Thus  $\mu_i := \text{rk } E_i$  is the multiplicity of the eigenvalue  $P_{ij}$  of  $A_j$  (provided that  $P_{ij} \neq P_{kj}$  for  $k \neq i$ ).

We see that  $\mu_0 = 1$ ,  $\sum_{i=0}^d \mu_i = n$ , and  $\mu_i = \text{tr } E_i = n(E_i)_{jj}$  (indeed,  $E_i$  has only eigenvalues 0 and 1, so  $\text{rk } E_i$  equals the sum of the eigenvalues).

**3.2. Theorem.** *The numbers  $P_{ij}$  and  $Q_{ij}$  satisfy*

- (i)  $P_{i0} = Q_{i0} = 1$ ,  $P_{0i} = n_i$ ,  $Q_{0i} = \mu_i$ ,
- (ii)  $P_{ij} P_{ik} = \sum_{l=0}^d p_{jk}^l P_{il}$ ,
- (iii)  $\mu_i P_{ij} = n_j Q_{ji}$ ,  $\sum_i \mu_i P_{ij} P_{ik} = nn_j \delta_{jk}$ ,  $\sum_i n_i Q_{ij} Q_{ik} = n\mu_j \delta_{jk}$ ,
- (iv)  $|P_{ij}| \leq n_j$ ,  $|Q_{ij}| \leq \mu_j$ .

**Proof.** Part (i) follows easily from  $\sum_i E_i = I = A_0$ ,  $\sum_i A_i = J = nE_0$ ,  $A_i J = n_i J$ , and  $\text{tr } E_i = \mu_i$ .

Part (ii) follows from  $A_j A_k = \sum_l p_{jk}^l A_l$ .

The first equality in (iii) follows from  $\sum_i n_j Q_{ji} P_{ik} = nn_j \delta_{jk} = \text{tr } A_j A_k = \sum_i \mu_i P_{ij} P_{ik}$ , since  $P$  is nonsingular,

this also proves the second equality, and the last one follows since  $PQ = nI$ .

The first inequality of (iv) holds because the  $P_{ij}$  are eigenvalues of the  $n_j$ -regular graphs  $(X, R_j)$ . The



second inequality then follows by use of (iii). ■

Relations (iii) are often referred to as the *orthogonality relations*, since they state that the rows (and columns) of  $P$  (and  $Q$ ) are orthogonal with respect to a suitable weight function.

If  $d = 2$ , and  $(X, R_1)$  is strongly regular with parameters  $(v, k, \lambda, \mu)$ , the matrices  $P$  and  $Q$  are

$$P = \begin{bmatrix} 1 & k & v-k-1 \\ 1 & r & -r-1 \\ 1 & s & -s-1 \end{bmatrix}, \quad Q = \begin{bmatrix} 1 & f & g \\ 1 & f \frac{r}{k} & g \frac{s}{k} \\ 1 & -f \frac{r+1}{v-k-1} & -g \frac{s+1}{v-k-1} \end{bmatrix},$$

where  $r, s, f$  and  $g$  can be expressed in terms of  $v, k, \lambda$  by use of (2.2).

In general the matrices  $P$  and  $Q$  can be computed from the intersection numbers of the scheme, as follows from the following

**3.3. Theorem.** For  $i = 0, \dots, d$ , the intersection matrix  $L_j$  has eigenvalues  $P_{ij}$  ( $0 \leq i \leq d$ ).

**Proof.** (3.2.ii) yields

$$P_{ij} \sum_k P_{ik} (P^{-1})_{km} = \sum_{k,l} P_{il} (L_j)_{lk} (P^{-1})_{km},$$

hence  $PL_j P^{-1} = \text{diag}(P_{0j}, \dots, P_{dj})$ . ■

Thanks to this theorem, it is relatively easy to compute  $P, Q (= \frac{1}{n} P^{-1})$  and  $\mu_i (= Q_{0i})$ . It is also possible to express  $P$  and  $Q$  in terms of the (common) eigenvectors of the  $L_j$ . Indeed,  $PL_j P^{-1} = \text{diag}(P_{0j}, \dots, P_{dj})$  implies that the rows of  $P$  are left eigenvectors and the columns of  $Q$  are right eigenvectors. In particular,  $\mu_i$  can be computed from the right eigenvector  $u_i$  and the left eigenvector  $v_i^T$ , normalized such that  $(u_i)_0 = (v_i)_0 = 1$ , by use of  $\mu_i u_i^T v_i = n$ . Clearly, each  $\mu_i$  must be an integer. These are the *rationality conditions* for an association scheme. As we saw in the case of a strongly regular graph, these conditions can be very powerful. Godsil (see his chapter) puts the rationality conditions in a more general form, which is not restricted to association schemes.

### The Krein parameters.

The Bose-Mesner algebra  $A$  is not only closed under ordinary matrix multiplication, but also under componentwise (Hadamard, Schur) multiplication (denoted  $\circ$ ). Clearly  $\{A_0, \dots, A_d\}$  is the basis of minimal idempotents with respect to this multiplication.

Write

$$E_i \circ E_j = \frac{1}{n} \sum_{k=0}^d q_{ij}^k E_k.$$

The numbers  $q_{ij}^k$  thus defined are called the *Krein parameters*. (Our  $q_{ij}^k$  are those of Delsarte, but differ from SEIDEL [49]'s by a factor  $n$ .) As expected, we now have the analogue of (3.1) and (3.2).

**3.4. Theorem.** The Krein parameters of an association scheme satisfy

- (i)  $q_{0j}^k = \delta_{jk}, \quad q_{ij}^0 = \delta_{ij} \mu_j, \quad q_{ij}^k = q_{ji}^k.$
- (ii)  $\sum_i q_{ij}^k = \mu_j, \quad \sum_j \mu_j = n,$
- (iii)  $q_{ij}^k \mu_k = q_{ik}^j \mu_j,$
- (iv)  $\sum_l q_{ij}^l q_{kl}^m = \sum_l q_{kj}^l q_{il}^m,$
- (v)  $Q_{ij} Q_{ik} = \sum_{l=0}^d q_{jk}^l Q_{il},$
- (vi)  $n \mu_k q_{ij}^k = \sum_l n_l Q_{il} Q_{lj} Q_{ik}.$

**Proof.** Let  $\Sigma(A)$  denote the sum of all entries of the matrix  $A$ . Then  $JAJ = \Sigma(A)J$ ,  $\Sigma(A \circ B) = \text{tr } AB^T$  and

$\Sigma(E_i) = 0$  if  $i \neq 0$ , since then  $E_i J = n E_i E_0 = 0$ . Now (i) follows by use of  $E_i \circ E_0 = \frac{1}{n} E_i$ ,  $q_{ij}^0 = \Sigma(E_i \circ E_j) = \text{tr } E_i E_j = \delta_{ij} \mu_j$ , and  $E_i \circ E_j = E_j \circ E_i$ , respectively. Equation (iv) follows by evaluating  $E_i \circ E_j \circ E_k$  in two ways, and (iii) follows from (iv) by taking  $m = 0$ . Equation (v) follows from evaluating  $A_i \circ E_j \circ E_k$  in two ways, and (vi) follows from (v), using the orthogonality relation  $\sum_l n_l Q_{lm} Q_{lk} = \delta_{mk} \mu_k n$ .

Finally, by use of (iii) we have

$$\mu_k \sum_j q_{ij}^k = \sum_j q_{ik}^j \mu_j = n \cdot \text{tr } (E_i \circ E_k) = n \sum_l (E_i)_{ll} (E_k)_{ll} = \mu_i \mu_k,$$

proving (ii). ■

The above results illustrate a dual behaviour between ordinary multiplication, the numbers  $p_{ij}^k$  and the matrices  $A_i$  and  $P$  on the one hand, and Schur multiplication, the numbers  $q_{ij}^k$  and the matrices  $E_i$  and  $Q$  on the other hand. If two association schemes have the property that the intersection numbers of one are the Krein parameters of the other, then the converse is also true. Two such schemes are said to be (formally) dual to each other. One scheme may have several (formal) duals, or none at all (but when the scheme is invariant under a regular abelian group, there is a natural way to define a dual scheme, cf. DELSARTE [17]). In fact usually the Krein parameters are not even integers. But they cannot be negative. These important restrictions, due to SCOTT [48], are the so-called Krein conditions.

**3.5. Theorem.** *The Krein parameters of an association scheme satisfy  $q_{ij}^k \geq 0$  for all  $i, j, k \in \{0, \dots, d\}$ .*

**Proof.** The numbers  $\frac{1}{n} q_{ij}^k$  ( $0 \leq k \leq d$ ) are the eigenvalues of  $E_i \circ E_j$  (since  $(E_i \circ E_j) E_k = \frac{1}{n} q_{ij}^k E_k$ ). On the other hand, the Kronecker product  $E_i \otimes E_j$  is positive semidefinite, since each  $E_i$  is. But  $E_i \circ E_j$  is a principal submatrix of  $E_i \otimes E_j$ , and therefore is positive semidefinite as well, i.e., has no negative eigenvalue. ■

The Krein parameters can be computed by use of equation (3.4.vi). This equation also shows that the Krein condition is equivalent to

$$\sum_l n_l Q_{li} Q_{lj} Q_{lk} \geq 0 \quad \text{for all } i, j, k \in \{0, \dots, d\}.$$

In case of a strongly regular graph we obtain

$$q_{11}^1 = \frac{f^2}{v} \left[ 1 + \frac{r^3}{k^2} - \frac{(r+1)^3}{(v-k-1)^2} \right] \geq 0,$$

$$q_{22}^2 = \frac{g^2}{v} \left[ 1 + \frac{s^3}{k^2} - \frac{(s+1)^3}{(v-k-1)^2} \right] \geq 0$$

(the other Krein conditions are trivially satisfied in this case), which is equivalent to the result mentioned in the previous section.

NEUMAIER [45], generalized Seidel's absolute bound to association schemes, and obtained

**3.6. Theorem.** *The multiplicities  $\mu_i$  ( $0 \leq i \leq d$ ) of an association scheme with  $d$  classes satisfy*

$$\sum_{q_{ij}^k \neq 0} \mu_k \leq \begin{cases} \mu_i \mu_j & \text{if } i \neq j, \\ \frac{1}{2} \mu_i (\mu_i + 1) & \text{if } i = j. \end{cases}$$

**Proof.** The left hand side equals  $\text{rk}(E_i \circ E_j)$ . But  $\text{rk}(E_i \circ E_j) \leq \text{rk}(E_i \otimes E_j) = \text{rk } E_i \times \text{rk } E_j = \mu_i \mu_j$ . And if  $i = j$ , then  $\text{rk}(E_i \circ E_i) \leq \frac{1}{2} \mu_i (\mu_i + 1)$ . Indeed, if the rows of  $E_i$  are linear combinations of  $\mu_i$  rows, then the rows of  $E_i \circ E_i$  are linear combinations of the  $\mu_i + \frac{1}{2} \mu_i (\mu_i - 1)$  rows that are the elementwise products of any two of these  $\mu_i$  rows. ■

For strongly regular graphs with  $q_{11}^1 = 0$  we obtain Seidel's bound:  $v \leq \frac{1}{2} f(f+3)$ . But in case  $q_{11}^1 > 0$ , Neumaier's result states that the bound can be improved to  $v \leq \frac{1}{2} f(f+1)$ .

### Subsets of association schemes.

The last subject of this section is a result of DELSARTE [17] (Theorem 3.3, p. 26) on subsets in association schemes. For  $Y \subseteq X$ ,  $Y \neq \emptyset$ , we define

$$a_i = \frac{1}{|Y|} \chi^T A_i \chi,$$

where  $\chi$  is the characteristic vector of  $Y$ . In other words,  $a_i$  is the average degree of the subgraph of  $(X, R_i)$  induced by  $Y$ . Clearly  $a_0 = 1$ , and  $\sum_{i=0}^d a_i = |Y|$ . The vector  $a = (a_0, \dots, a_d)$  is called the *inner distribution* of  $Y$ .

**3.7. Theorem.** *The inner distribution  $a$  of a nonempty subset of an association scheme satisfies  $aQ \geq 0$ .*

**Proof.**  $|Y| \sum_{i=0}^d a_i Q_{ij} = \sum_{i=0}^d \chi^T Q_{ij} A_i \chi = n \chi^T E_j \chi \geq 0$ , since  $E_j$  is positive semidefinite. ■

This inequality leads to Delsarte's linear programming bound, as we shall see in the next section.

As an application we have the following result (DELSARTE [17], Theorem 3.9, p. 32).

**3.8. Theorem.** *Let  $\{\{0\}, I_1, I_2\}$  be a partition of  $\{0, \dots, d\}$ , and assume that  $Y$  and  $Z$  are nonempty subsets of  $X$  such that the inner distribution  $b$  of  $Y$  satisfies  $b_i = 0$  for  $i \in I_1$ , and the inner distribution  $c$  of  $Z$  satisfies  $c_i = 0$  for  $i \in I_2$ . Then  $|Y| \cdot |Z| \leq |X|$ , and equality holds if and only if for all  $i \neq 0$  we have  $(bQ)_i = 0$  or  $(cQ)_i = 0$ .*

**Proof.** Define  $\beta_i = \mu_i^{-1} |Z|^{-1} \sum_j c_j Q_{ji}$ . Then  $\beta_0 = 1$ ,  $\beta_i \geq 0$  for all  $i$ , and  $\sum_i \beta_i Q_{ki} = n_k^{-1} |Z|^{-1} n c_k$ . Now we have  $|Y| = \sum_k b_k = (bQ)_0 \leq \sum_{i,k} b_k Q_{ki} \beta_i = \sum_i Q_{0i} \beta_i = \frac{n}{|Z|}$ . ■

Let us investigate a special case of this situation somewhat closer. Let, for  $I \subseteq \{0, \dots, d\}$ , the  $I$ -sphere around the point  $x \in X$  be the set  $\{y \in X | (x, y) \in R_i \text{ for some } i \in I\}$ . A nonempty subset  $Y$  of  $X$  is called *perfect* (more precisely, *I-perfect*) when the  $I$ -spheres around its points form a partition of  $X$ .

**3.9. Theorem.** ('Lloyd's theorem', cf. LLOYD [42], DELSARTE [17], p. 63). *Let  $Y$  be I-perfect, with inner distribution  $a$ . Then  $\sum_{i \in I} P_{ji} = 0$  for all  $j \neq 0$  such that  $(aQ)_j \neq 0$ .*

**Proof.** Apply the previous theorem, with for  $Z$  an  $I$ -sphere. If  $c$  is the inner distribution of  $Z$ , then

$$|Z| \mu_j^{-1} (cQ)_j = \mu_j^{-1} \sum_{i \in I} \sum_{g, h \in I} n_g P_{ih}^g Q_{ij} = \sum_{i \in I} \sum_{g, h \in I} P_{gh}^i P_{ji} = \sum_{g, h \in I} P_{jk} P_{jh} = \left( \sum_{i \in I} P_{ji} \right)^2. \quad \blacksquare$$

## 4. Applications.

In this section we discuss some special types of association scheme and their significance to other fields of combinatorics.

### Distance regular graphs.

Consider a connected simple graph with vertex set  $X$  of diameter  $d$ . Define  $R_i \subset X^2$  by  $(x, y) \in R_i$  whenever  $x$  and  $y$  have graph distance  $i$ . If this defines an association scheme, then the graph  $(X, R_1)$  is called *distance-regular*. The corresponding association scheme is called *metric*. By the triangle inequality,  $p_{ij}^k = 0$  if  $i + j < k$  or  $|i - j| > k$ . Moreover,  $p_{ij}^{i+j} > 0$ . Conversely, if the intersection numbers of an association scheme satisfy these conditions, then  $(X, R_1)$  is easily seen to be distance-regular.

Many of the association schemes that play a rôle in combinatorics are metric. In fact, all the examples treated in this chapter are metric. Strongly regular graphs are obviously metric. The line graph of the Petersen graph and the Hoffman-Singleton graph are easy examples of distance-regular graphs that are not strongly regular.

Any  $k$ -regular graph of diameter  $d$  has at most



$$1 + k + k(k-1) + \dots + k(k-1)^{d-1}$$

vertices, as is easily seen. Graphs for which equality holds are called *Moore graphs*. Moore graphs are distance-regular, and those of diameter 2 were dealt with in Theorem 2.3. Using the rationality conditions DAMERILL [16] and BANNAI & ITO [2] showed:

**4.1. Theorem.** A Moore graph with diameter  $d \geq 3$  is a  $(2d+1)$ -gon.

A strong non-existence result of the same nature is the theorem of FET & G. HIGMAN [22] about finite generalized polygons. A *generalized  $m$ -gon* is a point-line geometry such that the incidence graph is a connected, bipartite graph of diameter  $m$  and girth  $2m$ . It is called *regular* of order  $(s, t)$  for certain (finite or infinite) cardinal numbers  $s, t$  if each line is incident with  $s+1$  points and each point is incident with  $t+1$  lines. (It is not difficult to prove that if each point is on at least three lines, and each line has at least three points (and  $m < \infty$ ), then the geometry is necessarily regular, and in fact  $s = t$  in case  $m$  is odd.) From such a regular generalized  $m$ -gon of order  $(s, t)$ , where  $s$  and  $t$  are finite and  $m \geq 3$ , we can construct a distance-regular graph with valency  $s(t+1)$  and diameter  $d = \lceil \frac{1}{2}m \rceil$  by taking the collinearity graph on the points.

**4.2. Theorem.** A finite generalized  $m$ -gon of order  $(s, t)$  with  $s > 1$  and  $t > 1$  satisfies  $m \in \{2, 3, 4, 6, 8\}$ .

Proofs of this theorem can be found in FET & HIGMAN [22], KILMOYER & SOLOMON [36] and ROOS [47]; again the rationality conditions do the job. The Krein conditions yield some additional information:

**4.3. Theorem.** A finite regular generalized  $m$ -gon with  $s > 1$  and  $t > 1$  satisfies  $s \leq t^2$  and  $t \leq s^2$  if  $m = 4$  or  $8$ ; it satisfies  $s \leq t^3$  and  $t \leq s^3$  if  $m = 6$ .

This result is due to HIGMAN [28, 29] and HAEMERS & ROOS [26]. For each  $m \in \{2, 3, 4, 6, 8\}$  infinitely many generalized  $m$ -gons exist. (For  $m = 2$  we have trivial structures - the incidence graph is complete bipartite; for  $m = 3$  we have (generalized) projective planes; an example of a generalized 4-gon of order  $(2, 2)$  with collinearity graph  $T(6)$  can be described as follows: the points are the pairs from a 6-set, and the lines are the partitions of the 6-set into three pairs, with obvious incidence.)

Many association schemes have the important property that the eigenvalues  $P_{ij}$  can be expressed in terms of orthogonal polynomials. An association scheme is called  *$P$ -polynomial* if there exist polynomials  $f_k$  of degree  $k$  with real coefficients, and real numbers  $z_i$  such that  $P_{ik} = f_k(z_i)$ . Clearly we may always take  $z_i = P_{i1}$ .

By the orthogonality relation (3.2.iii) we have

$$\sum_i \mu_i f_j(z_i) f_k(z_i) = \sum_i \mu_i P_{ij} P_{ik} = n n_j \delta_{jk},$$

which shows that the  $f_k$  are orthogonal polynomials.

**4.4. Theorem.** An association scheme is metric if and only if it is  $P$ -polynomial.

**Proof.** Let the scheme be metric. Theorem 1.1 gives

$$A_1 A_i = p_{1i}^{i-1} A_{i-1} + p_{1i}^i A_i + p_{1i}^{i+1} A_{i+1}.$$

Since  $p_{1i}^{i+1} \neq 0$ ,  $A_{i+1}$  can be expressed in terms of  $A_1$ ,  $A_{i-1}$  and  $A_i$ . Hence for each  $j$  there exists a polynomial  $f_j$  of degree  $j$  such that

$$A_j = f_j(A_1).$$

Using this we have  $P_{ij} E_i = A_j E_i = f_j(A_1) E_i = f_j(A_1 E_i) E_i = f_j(P_{i1}) E_i$ , hence  $P_{ij} = f_j(P_{i1})$ .

Now suppose that the scheme is  $P$ -polynomial. Then the  $f_j$  are orthogonal polynomials, and therefore they satisfy a 3-term recurrence relation (see SZÉCÖ [52] p. 42)

$$\alpha_{j+1} f_{j+1}(z) = (\beta_j - z) f_j(z) + \gamma_{j-1} f_{j-1}(z).$$

Hence

$$P_{i1} P_{ij} = -\alpha_{j+1} P_{i,j+1} + \beta_j P_{ij} + \gamma_{j-1} P_{i,j-1} \quad \text{for } i = 0, \dots, d.$$

Since  $P_{i1} P_{ij} = \sum_l p_{1ij}^l P_{il}$  and  $P$  is nonsingular, it follows that  $p_{1ij}^l = 0$  for  $|l-j| > 1$ . Now the full metric property easily follows by induction. ■

This result is due to DELSARTE [17] (Theorem 5.6, p. 61). There is also a result dual to this theorem, involving so-called  $Q$ -polynomial and cometric schemes. However, just as the intersection numbers  $p_{ij}^k$  have a combinatorial interpretation while the Krein parameters  $q_{ij}^k$  do not, the metric schemes have the combinatorial description of distance-regular graphs, while there is no combinatorial interpretation for the cometric property. For more information on  $P$ - and  $Q$ -polynomial association schemes, see DELSARTE [17] and BANNAI & ITO [3]; for distance-regular graphs, see the forthcoming book by Brouwer, Cohen & Neumaier.

#### The Hamming scheme and error-correcting codes.

Let  $X = Q^d$ , the set of all vectors of length  $d$  with entries in  $Q$ , where  $Q$  is some set of size  $q$ . Define  $R_i \subset X^2$  by  $(x, y) \in R_i$  if the Hamming distance between  $x$  and  $y$  (i.e., the number of coordinates in which  $x$  and  $y$  differ) equals  $i$ . This defines an association scheme, the *Hamming scheme*  $H(d, q)$ . The Hamming scheme is easily seen to be metric, and hence by (4.4)  $P$ -polynomial. The orthogonal polynomials involved are the Kravčuk polynomials  $K_j(x)$ .

**4.5. Theorem.** *For the Hamming scheme  $H(d, q)$  we have*

$$P_{ij} = Q_{ij} = K_j(i) = \sum_{k=0}^j (-1)^k (q-1)^{j-k} \binom{i}{k} \binom{n-i}{j-k}.$$

See DELSARTE [17] (p. 38) or MACWILLIAMS & SLOANE [43] for proofs. From (4.5) we see that  $P = Q$ , so the Hamming scheme is self-dual.

A subset  $Y \subset X$  of  $H(d, q)$ , such that  $Y^2 \cap R_i = \emptyset$  for  $i = 1, \dots, \delta-1$  and  $Y^2 \cap R_\delta \neq \emptyset$  (i.e., a subset  $Y$  such that the minimum Hamming distance between two vectors of  $Y$  equals  $\delta$ ), is nothing but an error-correcting code with parameters  $(d, |Y|, \delta)$  over the alphabet  $Q$  (cf. the chapter of coding theory). Let  $a = (a_0, \dots, a_d)$  be the inner distribution of  $Y$ . Then  $\sum_i a_i = |Y|$ ,  $a_0 = 1$ ,  $a_1 = \dots = a_{\delta-1} = 0$ ,  $a_i \geq 0$  (by definition), and  $aQ \geq 0$  by (3.7). Consider  $a_\delta, \dots, a_d$  as variables and define

$$a^* = 1 + \max \sum_{i=\delta}^d a_i \text{ subject to } K_j(1) + \sum_{i=\delta}^d a_i K_j(i) \geq 0, j = 0, \dots, d, \text{ and } a_i \geq 0, i = \delta, \dots, d.$$

Then clearly  $|Y| \leq a^*$ . So  $a^*$  is an upper bound for the number of codewords with a given length and minimum distance. This bound, due to DELSARTE [17], is called the linear programming bound, since the value of  $a^*$  can be computed by linear programming. Of course, the above-mentioned inequalities are not the only ones satisfied by the  $a_i$ , and by adding extra inequalities to the system, one may obtain sharper bounds on  $|Y|$ . For details and more applications to coding theory, see DELSARTE [17], MACWILLIAMS & SLOANE [43], BEST, BROUWER, MACWILLIAMS, ODLYZKO & SLOANE [6], BEST & BROUWER [5].

#### The Johnson scheme and $t$ -designs.

Let the set  $X$  consist of all subsets of size  $d$  of a set  $M$ , where  $|M| = m \geq 2d$ . Define relation  $R_i \subset X^2$  by  $(x, y) \in R_i$  if the Johnson distance between  $x$  and  $y$  (i.e., the cardinality of  $x \setminus y$ ) equals  $i$ . This defines an association scheme, the *Johnson scheme*  $J(d, m)$ . Since the Johnson distance between  $x$  and  $y$  equals twice the Hamming distance between (the characteristic vectors of)  $x$  and  $y$ , it follows that also the Johnson distance satisfies the triangle inequality, so that the Johnson scheme is metric. Note that the graph  $(X, R_1)$  is complete for  $d = 1$ , and is the triangular graph  $T(m)$  for  $d = 2$ . The following result (due to OGASAWARA [46] and YAMAMOTO et al. [54]) gives some parameters:

**4.6. Theorem.** *For the Johnson scheme  $J(d, m)$  the following hold:*

$$P_{ij} = \frac{n_j}{\mu_i} Q_{ji} = E_j(i) = \sum_{k=0}^j (-1)^k \binom{d-k}{j-k} \binom{d-i}{k} \binom{m-d+k-i}{k} = \sum_{k=0}^j (-1)^k \binom{i}{k} \binom{d-i}{j-k} \binom{m-d-i}{j-k},$$

$$\mu_i = \binom{m}{i} - \binom{m}{i-1}, \quad n_j = \binom{d}{j} \binom{m-d}{j}.$$

Here  $E_j(x)$  is a so-called Eberlein polynomial. It has degree  $2j$  in the indeterminate  $x$ , and degree  $j$  in the indeterminate  $x(m+1-x)$ . Since  $P_{i1} = d(m-1) - i(m+1-i)$ ,  $E_j(i)$  indeed has degree  $j$  in  $P_{i1}$  as required by the definition of  $P$ -polynomial scheme.



The graph  $(X, R_d)$  of a Johnson scheme is called a Kneser graph. A subset  $Y$  of  $X$  such that any two elements of  $Y$  have non-empty intersection is a coclique (independent set) of the Kneser graph. By use of Theorem 3.7 and the above formulas, it can be deduced that

$$|Y| \leq \binom{m-1}{d-1},$$

the famous result of ERDŐS, KO & RADO [21].

A  $t$ -( $m, d, \lambda$ ) design is a subset  $Y \subset X$  of the Johnson scheme  $J(d, m)$  such that each  $t$ -element subset of  $M$  is contained in precisely  $\lambda$  elements of  $Y$ . DELSARTE [17] (Theorem 4.7, p. 51) proved the following:

**4.7. Theorem.** A non-empty subset  $Y$  of the Johnson scheme  $J(d, m)$  with inner distribution  $a = (a_0, \dots, a_d)$  is a  $t$ -( $m, d, \lambda$ ) design if and only if

$$\sum_{i=0}^d a_i Q_{ij} = 0 \quad \text{for } j = 1, \dots, t. \quad (*)$$

Just as we did in case of the Hamming scheme, we can define

$$a_* = 1 + \min \sum_{i=1}^d a_i \quad \text{subject to} \\ a_i \geq 0 \text{ for } i = 1, \dots, d, \text{ and } \sum_{i=0}^d a_i Q_{ij} = 0 \text{ for } j = 1, \dots, t, \text{ and } \sum_{i=0}^d a_i Q_{ij} \geq 0 \text{ for } j = t+1, \dots, d,$$

where the  $Q_{ij}$  are found in (4.6). Now we have the linear programming (lower) bound for the number of blocks in a  $t$ -design:  $|Y| \geq a_*$ . Using the simplex algorithm, HAEMERS & WEUG [27] showed that this inequality implies the non-existence of the designs with parameters 4-(17, 8, 5), 4-(23, 11,  $\lambda$ ),  $\lambda = 6, 12$ , 4-(24, 12, 15), 6-(19, 9,  $\lambda$ ),  $\lambda \leq 10$ , 6-(20, 10,  $\lambda$ ),  $\lambda = 7, 14$ . In certain other cases, such as 5-(19, 9, 7), KÖHLER [34] ruled out the existence of  $t$ -designs by showing that no solution of the above system of inequalities corresponds to an actual design.

If we replace the restrictions  $(aQ)_i = 0$  by restrictions  $a_i = 0$  ( $1 \leq i < \frac{1}{2}\delta$ ) in this system of inequalities, and maximize  $\sum a_i$ , we obtain upper bounds for the cardinality of codes with minimum distance  $\delta$  and constant weight  $d$ .

We might also require both  $a_i = 0$  and  $(aQ)_j = 0$  for suitable  $i, j$ , and obtain results for  $t$ -designs with restricted block intersections, such as quasisymmetric block designs. By this method it follows for instance that a quasisymmetric 2-(29, 7, 12) design with block intersections 1 and 3 cannot exist (HAEMERS [24]). For details and more results we refer to DELSARTE [17], MACWILLIAMS & SLOANE [43], BEST, BROUWER, MACWILLIAMS, ODLYZKO & SLOANE [6], CAMERON & VAN LINT [15], CALDERBANK [13]. We also point out that several results of §3 in Godsil's chapter can also be obtained by use of the framework of Johnson schemes.

DELSARTE [17] generalized the notion of  $t$ -designs to subsets of arbitrary association schemes satisfying (\*). (Equivalently, a  $t$ -design is a subset  $Y$  of  $X$  such that its characteristic vector  $\chi$  satisfies  $E_j \chi = 0$  for  $j = 1, \dots, t$ .) He shows that a  $t$ -design in the Hamming scheme is what is known as an orthogonal array of strength  $t$  (see the chapter on Block Designs). Thus, we also have a linear programming bound for orthogonal arrays. More generally, one may give an interpretation of the classical concept of  $t$ -design in terms of ranked posets in the obvious way, and then prove for each of the eight known infinite families of  $P$ - and  $Q$ -polynomial association schemes that a subset is a classical  $t$ -design if and only if it is a Delsarte  $f(t)$ -design (where usually  $f(t) = t$ ), see DELSARTE [19] and STANTON [51].

### Imprimitive schemes.

In section 2 we have seen a completely different relation between designs and association schemes. Let us give one more example. Let  $N$  be the incidence matrix of a square 2-design. Then, defining  $A_0 = I$ ,

$$A_1 = \begin{bmatrix} 0 & N \\ N^T & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} J & 0 \\ 0 & J \end{bmatrix}$$

and  $A_3 = J - I - A_1 - A_2$ , we obtain a 3-class association scheme. It is *imprimitive*, that is, the union of some of the  $R_i$  form a non-trivial equivalence relation (here  $R_2$  is an equivalence relation). Another

imprimitive association scheme we have seen is the line graph of the Petersen graph (there having maximal distance is an equivalence relation).

Given an imprimitive association scheme one may produce new association schemes; on the one hand, there is a natural way to give the set of equivalence classes the structure of an association scheme (the 'quotient scheme'), and on the other hand, each equivalence class together with the restrictions of the original relations becomes an association scheme (a 'subscheme' of the original scheme).

### The group case.

We very briefly discuss some relations between association schemes and finite permutation groups.

Let  $G$  be a permutation group acting on a set  $X$ . Then  $G$  has a natural action on  $X^2$ , the orbits of which are called orbitals. Suppose  $G$  acts generously transitive on  $X$ , that is, for any  $x, y \in X$  there exists an element of  $G$  interchanging  $x$  and  $y$ . Then the orbitals form an association scheme.

(Without any requirements on  $G$ , the orbitals form a coherent configuration (see § 3). The coherent configuration is homogeneous if  $G$  is transitive. We get an association scheme in the sense of Delsarte when the permutation character is multiplicity free.)

For any  $x \in X$ , the number of orbitals equals the number of orbits on  $X$  of  $G_x$  (the subgroup of  $G$  of permutations fixing  $x$ ). This number is called the *rank* of  $G$ . Thus, the number of classes in the association scheme is one less than the rank of  $G$ . We can also transfer other permutation group theoretic terminology and results to the theory of association schemes. For instance, the Bose-Mesner algebra is in the group case known as the *centralizer algebra*, and all standard results on this centralizer algebra (cf., e.g., WIELANDT [53]) have their direct analogue for the Bose-Mesner algebra.

The Hamming and Johnson schemes are derived from generously transitive permutation representations as discussed above; for instance, the Johnson scheme is derived from the representation of the symmetric group  $\text{Sym}(m)$  on the  $d$ -element subsets of an  $m$ -set.

If a metric association scheme belongs to the group case, then the corresponding distance-regular graph is called *distance-transitive*. In other words, a graph is distance-transitive when its group of automorphisms is transitive on pairs of vertices with a given distance. Distance-transitive strongly regular graphs are known as rank 3 graphs. A rank 3 permutation group is generously transitive if and only if it has even order; consequently every rank 3 permutation group of even order provides a strongly regular graph. All such strongly regular graphs have recently been classified, see KANTOR & LIEBLER [35], LIEBECK [38, 40], and LIEBECK & SAXL [39].

### References

1. Aschbacher, M., *The non-existence of rank three permutation groups of degree 3250 and subdegree 57*, J. Algebra **19** (1971) 538-540.
2. Bannai, E. and T. Ito, *On finite Moore graphs*, J. Fac. Sci. Univ. Tokyo, Sect. IA **20** (1973) 191-208.
3. Bannai, E. and T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin-Cummings Lecture Note Ser. 58, The Benjamin/Cummings Publishing Company, Inc., London (1984).
4. Belevitch, V., *Theory of 2n-terminal networks with application to conference telephony*, Elect. Commun. **27** (1950) 231-244.
5. Best, M.R. and A.E. Brouwer, *The triply shortened binary Hamming code is optimal*, Discr. Math. **17** (1977) 235-245.
6. Best, M.R., A.E. Brouwer, F.J. MacWilliams, A.M. Odlyzko, and N.J.A. Sloane, *Bounds on binary codes of word length less than 25*, IEEE Trans. Inf. Theory **IT-24** (1978) 81-93.
7. Biggs, N.L., *Algebraic Graph Theory*, Cambridge Tracts in Math. 67, Cambridge University Press, Cambridge (1974).



8. Bose, R.C., *Strongly regular graphs, partial geometries, and partially balanced designs*, Pacific J. Math. **13** (1963) 389-419.
9. Bose, R.C. and D.M. Mesner, *On linear associative algebras corresponding to association schemes of partially balanced designs*, Ann. Math. Stat. **30** (1959) 21-38.
10. Bose, R.C. and T. Shimamoto, *Classification and analysis of partially balanced incomplete block designs with two associate classes*, J. Amer. Statist. Assoc. **47** (1952) 151-184.
11. Brouwer, A.E. and J.H. van Lint, *Strongly regular graphs and partial geometries*, pp. 85-122 in: *Enumeration and Design - Proc. Silver Jubilee Conf. on Combinatorics, Waterloo, 1982* (ed. D.M. Jackson & S.A. Vanstone), Academic Press, Toronto, 1984.
12. Burrow, M., *Representation theory of finite groups*, Academic Press, New York (1965).
13. Calderbank, A.R., *Geometric invariants for quasi-symmetric designs*, to appear.
14. Cameron, P.J., *Strongly regular graphs*, pp. 337-360 in: *Selected topics in graph theory* (ed. L.W. Beineke & R.J. Wilson), Acad. Press, 1978.
15. Cameron, P.J. and J.H. van Lint, *Graphs, Codes and Designs*, London Math. Soc. Lecture Notes 43, Cambridge Univ. Press, Cambridge (1980).
16. Damerell, R.M., *On Moore graphs*, Proc. Cambridge Phil. Soc. **74** (1973) 227-236.
17. Delsarte, Ph., *An algebraic approach to the association schemes of coding theory*, Philips Research Reports Suppl. **10** (1973).
18. Delsarte, Ph., *The association schemes of coding theory*, pp. 143-161 in: *Combinatorics* (ed. M. Hall, jr. & J.H. van Lint), Reidel, Dordrecht, 1975.
19. Delsarte, Ph., *Association schemes and t-designs in regular semilattices*, J. Combinatorial Th. (A) **20** (1976) 230-243.
20. Delsarte, Ph., J.-M. Goethals, and J.J. Seidel, *Bounds for systems of lines and Jacobi polynomials*, Philips Res. Reports **30** (1975) 91\*-105\*.
21. Erdős, P., Chao Ko, and R. Rado, *Intersection theorems for systems of finite sets*, Quart. J. Math. Oxford (2) **12** (1961) 313-320.
22. Feit, W. and G. Higman, *The non-existence of certain generalized polygons*, J. Algebra **1** (1964) 114-131.
23. Goethals, J.-M., *Association schemes*, pp. 157-180 in: *Algebraic coding theory and applications*, CISM courses and lectures 258 (ed. G. Longo), 1979.
24. Haemers, W.H., *Sterke grafen en block designs*, Master thesis, Eindhoven Univ. of Technology (1975). (Dutch)
25. Haemers, W.H., *Eigenvalue methods*, pp. 15-38 in: *Packing and covering in combinatorics*, Math. Centr. Tracts 106 (ed. A. Schrijver), Math. Centr., Amsterdam, 1979.
26. Haemers, W.H. and C. Roos, *An inequality for generalized hexagons*, Geometriae Dedicata **10** (1981) 219-222.
27. Haemers, W.H. and C. Weug, *Linear programming bounds for codes and designs*, Technical Note N96, M.B.L.E. Brussels (1974).
28. Higman, D.G., *Partial geometries, generalized quadrangles and strongly regular graphs*, pp. 263-293 in: *Atti del Convegno di Geometria Combinatoria e sue Applicazioni* (Univ. degli Studi di Perugia, Perugia, 1970), Perugia, 1971.
29. Higman, D.G., *Invariant relations, coherent configurations and generalized polygons*, pp. 27-43 in: *Combinatorics*, Math. Centre Tracts 57 (ed. M. Hall & J.H. van Lint), Amsterdam, 1974. (Reidel, Dordrecht)
30. Higman, D.G., *Coherent configurations, Part I: Ordinary representation theory*, Geometriae Dedicata **4** (1975) 1-32.

31. Higman, D.G., *Coherent configurations, Part II: Weights*, Geometriae Dedicata **5** (1976) 413-424.
32. Hoffman, A.J. and R.R. Singleton, *On Moore graphs with diameters 2 and 3*, IBM J. Res. Developm. **4** (1960) 497-504.
33. Hubaut, X., *Strongly regular graphs*, Discrete Math. **13** (1975) 357-381.
34. Köhler, E., *Über den allgemeinen Gray-Code und die Nichtexistenz einiger  $t$ -designs*, pp. 102-111 in: *Graphen in Forschung und Unterricht, Festschrift K. Wagner* (ed. Rainer Bodendiek), Franzbecker, Salzdetfurth, 1985.
35. Kantor, W.M. and R.A. Liebler, *The rank three representations of the finite classical groups*, Trans. Amer. Math. Soc. **271** (1982) 1-71.
36. Kilmoyer, R. and L. Solomon, *On the theorem of Feit-Higman*, J. Combinatorial Th. (A) **15** (1973) 310-322.
37. Koornwinder, T.H., *A note on the absolute bound for systems of lines*, Proc. KNAW A **79** (1976) 152-153. (= Indag. Math. **38**)
38. Liebeck, M.W., *Permutation modules for rank 3 symplectic and orthogonal groups*, J. Algebra **92** (1985) 9-15.
39. Liebeck, M.W. and J. Saxl, *The finite primitive permutation groups of rank three*, Bull. London Math. Soc. **18** (1986) 165-172.
40. Liebeck, M.W., *The affine permutation groups of rank three*, to appear.
41. Lint, J.H. van and J.J. Seidel, *Equilateral point sets in elliptic geometry*, Proc. Kon. Ned. Acad. Wet. A **69** (1966) 335-348. (= Indag. Math. **28**)
42. Lloyd, S.P., *Binary block coding*, Bell System Tech. J. **36** (1957) 517-535.
43. MacWilliams, F.J. and N.J.A. Sloane, *The Theory of Error-Correcting Codes*, North Holland Publ. Co., Amsterdam (1977).
44. Marcus, M. and H. Minc, *Introduction to linear algebra*, MacMillan, New York (1965).
45. Neumaier, A., *New inequalities for the parameters of an association scheme*, pp. 365-367 in: *Combinatorics and Graph Theory*, Springer Lecture Notes in Math. **885**, 1981.
46. Ogasawara, M., *A necessary condition for the existence of regular and symmetrical PBIB designs of  $T_m$  type*, Univ. North Carolina Inst. Statistics (1965).
47. Roos, C., *An alternative proof of the Feit-Higman theorem on generalized polygons*, Delft Progress Report **5** (1980) 67-77.
48. Scott, L.L. jr., *A condition on Higman's parameters*, Notices Amer. Math. Soc. **20** (1973) A-97.
49. Seidel, J.J., *Strongly regular graphs*, pp. 157-180 in: *Surveys in Combinatorics, Proc. 7th British Combinatorial Conference*, London Math. Soc. Lecture Note Ser. **38** (ed. B. Bollobás), Cambridge University Press, Cambridge, 1979.
50. Seidel, J.J., *Delsarte's theory of association schemes*, pp. 249-258 in: *Graphs and other combinatorial topics*, Teubner texte zur Mathematik **59** (ed. M. Fiedler), Leipzig, 1983.
51. Stanton, D.,  *$t$ -designs in Classical Association Schemes*, Graphs and Combinatorics **2** (1986) 283-286.
52. Szegő, G., *Orthogonal Polynomials*, Amer. Math. Soc. Colloquium Publ. **23**, AMS, New York (1959). (1st ed., 1939; 4th ed., Providence, 1975)
53. Wielandt, H., *Finite permutation groups*, Academic Press, New York (1964).
54. Yamamoto, S., Y. Fujii, and N. Hamada, *Composition of some series of association algebras*, J. Sci. Hiroshima Univ. Ser. A-I **29** (1965) 181-215.



## IN 1986 REEDS VERSCHENEN

- 202 J.H.F. Schilderincx  
Interregional Structure of the European Community. Part III
- 203 Antoon van den Elzen and Dolf Talman  
A new strategy-adjustment process for computing a Nash equilibrium in a noncooperative more-person game
- 204 Jan Vingerhoets  
Fabrication of copper and copper semis in developing countries. A review of evidence and opportunities
- 205 R. Heuts, J. van Lieshout, K. Baken  
An inventory model: what is the influence of the shape of the lead time demand distribution?
- 206 A. van Soest, P. Kooreman  
A Microeconomic Analysis of Vacation Behavior
- 207 F. Boekema, A. Nagelkerke  
Labour Relations, Networks, Job-creation and Regional Development. A view to the consequences of technological change
- 208 R. Alessie, A. Kapteyn  
Habit Formation and Interdependent Preferences in the Almost Ideal Demand System
- 209 T. Wansbeek, A. Kapteyn  
Estimation of the error components model with incomplete panels
- 210 A.L. Hempenius  
The relation between dividends and profits
- 211 J. Kriens, J.Th. van Lieshout  
A generalisation and some properties of Markowitz' portfolio selection method
- 212 Jack P.C. Kleijnen and Charles R. Standridge  
Experimental design and regression analysis in simulation: an FMS case study
- 213 T.M. Doup, A.H. van den Elzen and A.J.J. Talman  
Simplicial algorithms for solving the non-linear complementarity problem on the simplotope
- 214 A.J.W. van de Gevel  
The theory of wage differentials: a correction
- 215 J.P.C. Kleijnen, W. van Groenendaal  
Regression analysis of factorial designs with sequential replication
- 216 T.E. Nijman and F.C. Palm  
Consistent estimation of rational expectations models

- 217 P.M. Kort  
The firm's investment policy under a concave adjustment cost function
- 218 J.P.C. Kleijnen  
Decision Support Systems (DSS), en de kleren van de keizer ...
- 219 T.M. Doup and A.J.J. Talman  
A continuous deformation algorithm on the product space of unit simplices
- 220 T.M. Doup and A.J.J. Talman  
The 2-ray algorithm for solving equilibrium problems on the unit simplex
- 221 Th. van de Klundert, P. Peters  
Price Inertia in a Macroeconomic Model of Monopolistic Competition
- 222 Christian Mulder  
Testing Korteweg's rational expectations model for a small open economy
- 223 A.C. Meijdam, J.E.J. Plasmans  
Maximum Likelihood Estimation of Econometric Models with Rational Expectations of Current Endogenous Variables
- 224 Arie Kapteyn, Peter Kooreman, Arthur van Soest  
Non-convex budget sets, institutional constraints and imposition of concavity in a flexible household labor supply model
- 225 R.J. de Groof  
Internationale coördinatie van economische politiek in een twee-regio-twee-sectoren model
- 226 Arthur van Soest, Peter Kooreman  
Comment on 'Microeconomic Demand Systems with Binding Non-Negativity Constraints: The Dual Approach'
- 227 A.J.J. Talman and Y. Yamamoto  
A globally convergent simplicial algorithm for stationary point problems on polytopes
- 228 Jack P.C. Kleijnen, Peter C.A. Karremans, Wim K. Oortwijn, Willem J.H. van Groenendaal  
Jackknifing estimated weighted least squares
- 229 A.H. van den Elzen and G. van der Laan  
A price adjustment for an economy with a block-diagonal pattern
- 230 M.H.C. Paardekooper  
Jacobi-type algorithms for eigenvalues on vector- and parallel computer
- 231 J.P.C. Kleijnen  
Analyzing simulation experiments with common random numbers

- 232 A.B.T.M. van Schaik, R.J. Mulder  
On Superimposed Recurrent Cycles
- 233 M.H.C. Paardekooper  
Sameh's parallel eigenvalue algorithm revisited
- 234 Pieter H.M. Ruys and Ton J.A. Storcken  
Preferences revealed by the choice of friends
- 235 C.J.J. Huys en E.N. Kertzman  
Effectieve belastingtarieven en kapitaalkosten
- 236 A.M.H. Gerards  
An extension of König's theorem to graphs with no odd- $K_4$
- 237 A.M.H. Gerards and A. Schrijver  
Signed Graphs - Regular Matroids - Grafts
- 238 Rob J.M. Alessie and Arie Kapteyn  
Consumption, Savings and Demography
- 239 A.J. van Reeken  
Begrippen rondom "kwaliteit"
- 240 Th.E. Nijman and F.C. Palmer  
Efficiency gains due to using missing data. Procedures in regression models
- 241 Dr. S.C.W. Eijffinger  
The determinants of the currencies within the European Monetary System

## IN 1987 REEDS VERSCHENEN

- 242 Gerard van den Berg  
Nonstationarity in job search theory
- 243 Annie Cuyt, Brigitte Verdonk  
Block-tridiagonal linear systems and branched continued fractions
- 244 J.C. de Vos, W. Vervaat  
Local Times of Bernoulli Walk
- 245 Arie Kapteyn, Peter Kooreman, Rob Willemse  
Some methodological issues in the implementation  
of subjective poverty definitions
- 246 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel  
Sampling for Quality Inspection and Correction: AOQL Performance  
Criteria
- 247 D.B.J. Schouten  
Algemene theorie van de internationale conjuncturele en structurele  
afhankelijkheden
- 248 F.C. Bussemaker, W.H. Haemers, J.J. Seidel, E. Spence  
On  $(v,k,\lambda)$  graphs and designs with trivial automorphism group
- 249 Peter M. Kort  
The Influence of a Stochastic Environment on the Firm's Optimal Dyna-  
mic Investment Policy
- 250 R.H.J.M. Gradus  
Preliminary version  
The reaction of the firm on governmental policy: a game-theoretical  
approach
- 251 J.G. de Gooijer, R.M.J. Heuts  
Higher order moments of bilinear time series processes with symmetri-  
cally distributed errors
- 252 P.H. Stevers, P.A.M. Versteijne  
Evaluatie van marketing-activiteiten
- 253 H.P.A. Mulders, A.J. van Reeken  
DATAAL - een hulpmiddel voor onderhoud van gegevensverzamelingen
- 254 P. Kooreman, A. Kapteyn  
On the identifiability of household production functions with joint  
products: A comment
- 255 B. van Riel  
Was er een profit-squeeze in de Nederlandse industrie?
- 256 R.P. Gilles  
Economies with coalitional structures and core-like equilibrium con-  
cepts

- 257 P.H.M. Ruys, G. van der Laan  
Computation of an industrial equilibrium

Bibliotheek K. U. Brabant



17 000 01059992 7